

Package: cpd (via r-universe)

September 17, 2024

Type Package

Title Complex Pearson Distributions

Version 0.3.2

Date 2023-09-20

Maintainer Silverio Vilchez-Lopez <svilchez@ujaen.es>

Description Probability mass function, distribution function, quantile function and random generation for the Complex Triparametric Pearson (CTP) and Complex Biparametric Pearson (CBP) distributions developed by Rodriguez-Avi et al (2003) <[doi:10.1007/s00362-002-0134-7](https://doi.org/10.1007/s00362-002-0134-7)>, Rodriguez-Avi et al (2004) <[doi:10.1007/BF02778271](https://doi.org/10.1007/BF02778271)> and Olmo-Jimenez et al (2018) <[doi:10.1080/00949655.2018.1482897](https://doi.org/10.1080/00949655.2018.1482897)>. The package also contains maximum-likelihood fitting functions for these models.

Depends R (>= 4.0.0)

Imports hypergeo, Rdpack, dgof, graphics

RdMacros Rdpack

License GPL (>= 2)

Encoding UTF-8

RoxygenNote 7.2.1

NeedsCompilation no

Author Silverio Vilchez-Lopez [aut, cre], Maria Jose Olmo-Jimenez [aut], Jose Rodriguez-Avi [aut]

Date/Publication 2023-09-23 18:20:02 UTC

Repository <https://svilchez.r-universe.dev>

RemoteUrl <https://github.com/cran/cpd>

RemoteRef HEAD

RemoteSha af2df3d41d2f873c1e052400c4de2b285e7e9d4a

Contents

cbp	2
chisq.test2	4
ctp	5
ebw	7
fitcbp	9
fitctp	11
fitebw	13
plot.fitCBP	15
plot.fitCTP	16
plot.fitEBW	16
varcomp	17
Index	19

cbp *The Complex Biparametric Pearson (CBP) Distribution*

Description

Probability mass function, distribution function, quantile function and random generation for the Complex Biparametric Pearson (CBP) distribution with parameters b and γ .

Usage

```
dcbp(x, b, gamma)
pcbp(q, b, gamma, lower.tail = TRUE)
qcbp(p, b, gamma, lower.tail = TRUE)
rcbp(n, b, gamma)
```

Arguments

x	vector of (non-negative integer) quantiles.
b	parameter b (real)
gamma	parameter gamma (positive)
q	vector of quantiles.
lower.tail	if TRUE (default), probabilities are $P(X \leq x)$, otherwise, $P(X > x)$.
p	vector of probabilities.
n	number of observations. If $\text{length}(n) > 1$, the length is taken to be the number required.

Details

The CBP distribution with parameters b and γ has pmf

$$f(x|b, \gamma) = C\Gamma(ib + x)\Gamma(-ib + x)/(\Gamma(\gamma + x)x!), x = 0, 1, 2, \dots$$

where i is the imaginary unit, $\Gamma()$ the gamma function and

$$C = \Gamma(\gamma - ib)\Gamma(\gamma + ib)/(\Gamma(\gamma)\Gamma(ib)\Gamma(-ib))$$

the normalizing constant.

The CBP is a particular case of the CTP when $a = 0$.

The mean and the variance of the CBP distribution are $E(X) = \mu = b^2/(\gamma - 1)$ and $Var(X) = \mu(\mu + \gamma - 1)/(\gamma - 2)$ so $\gamma > 2$.

It is always overdispersed.

Value

dcbp gives the pmf, pcbp gives the distribution function, qcbp gives the quantile function and rcbp generates random values.

References

Jose Rodriguez-Avi J, Conde-Sanchez A, Saez-Castillo AJ (2003). "A new class of discrete distributions with complex parameters." *Stat. Pap.*, **44**, 67–88. doi:10.1007/s0036200201347.

See Also

Probability mass function, distribution function, quantile function and random generation for the CTP distribution: [dctp](#), [pctp](#), [qctp](#) and [rctp](#). Functions for maximum-likelihood fitting of the CBP distribution: [fitcbp](#).

Examples

```
# Examples for the function dcbp
dcbp(3,2,5)
dcbp(c(3,4),2,5)

# Examples for the function pcbp
pcbp(3,2,3)
pcbp(c(3,4),2,3)

# Examples for the function qcbp
qcbp(0.5,2,3)
qcbp(c(.8,.9),2,3)

# Examples for the function rcbp
rcbp(10,1,3)
```

 chisq.test2

Pearson's Chi-squared Test for Count Data

Description

chisq.test2 performs Pearson chi-squared goodness-of-fit test for count data

Usage

```
chisq.test2(obs, p.esp, npar = NULL, grouping = FALSE)
```

Arguments

obs	a numeric vector with the counts
p.esp	a numeric vector with the expected probabilities of the same length of obs. They must sum 1.
npar	an integer specifying the number of parameters of the model. By default npar is NULL, so the degrees of freedom of the chi-squared statistics are the number of classes minus 1.
grouping	a logical indicating whether to group in classes with expected frequency greater than or equal to 5. By default grouping is FALSE.

Value

A list with class "htest" containing the following components:

- `statistic`: the value of the chi-squared test statistic.
- `df`: the degrees of freedom of the approximate chi-squared distribution.
- `p.value`: the p-value for the test.
- `observed`: the observed counts.
- `observed.grouped`: the observed counts grouped in classes with expected frequency greater or equal to 5.
- `expected`: the expected counts under the null hypothesis.
- `expected.grouped`: the expected counts under the null hypothesis grouped in classes with expected frequency greater or equal to 5.
- `residuals`: the Pearson residuals computed as $(\text{observed} - \text{expected}) / \sqrt{\text{expected}}$.

Examples

```
set.seed(123)
x <- rctp(500, -1.5, 1, 2)
table(x)
obs <- c(172, 264, 57, 6, 0, 1)
fit <- fitctp(x)
p.esp <- c(dctp(0:(length(obs)-1), fit$coefficients[1], fit$coefficients[2]),
```

```

fit$coefficients[3])[1:(length(obs)-1)], 1-sum(dctp(0:(length(obs)-1),
fit$coefficients[1], fit$coefficients[2], fit$coefficients[3])[1:(length(obs)-1)]))
chisq.test2(obs, p.esp)
chisq.test2(obs, p.esp, grouping = TRUE)
chisq.test2(obs, npar= 3, p.esp)

```

ctp

The Complex Triparametric Pearson (CTP) Distribution

Description

Probability mass function, distribution function, quantile function and random generation for the Complex Triparametric Pearson (CTP) distribution with parameters a , b and γ .

Usage

```

dctp(x, a, b, gamma)

pctp(q, a, b, gamma, lower.tail = TRUE)

qctp(p, a, b, gamma, lower.tail = TRUE)

rctp(n, a, b, gamma)

```

Arguments

x	vector of (non-negative integer) quantiles.
a	parameter a (real)
b	parameter b (real)
gamma	parameter γ (positive)
q	vector of quantiles.
lower.tail	if TRUE (default), probabilities are $P(X \leq x)$, otherwise, $P(X > x)$.
p	vector of probabilities.
n	number of observations. If $\text{length}(n) > 1$, the length is taken to be the number required.

Details

The CTP distribution with parameters a , b and γ has pmf

$$f(x|a, b, \gamma) = C\Gamma(a + ib + x)\Gamma(a - ib + x)/(\Gamma(\gamma + x)x!), x = 0, 1, 2, \dots$$

where i is the imaginary unit, $\Gamma()$ the gamma function and

$$C = \Gamma(\gamma - a - ib)\Gamma(\gamma - a + ib)/(\Gamma(\gamma - 2a)\Gamma(a + ib)\Gamma(a - ib))$$

the normalizing constant.

If $a = 0$ the CTP is a Complex Biparametric Pearson (CBP) distribution, so the pmf of the CBP distribution is obtained. If $b = 0$ the CTP is an Extended Biparametric Waring (EBW) distribution, so the pmf of the EBW distribution is obtained.

The mean and the variance of the CTP distribution are $E(X) = \mu = (a^2 + b^2)/(\gamma - 2a - 1)$ and $Var(X) = \mu(\mu + \gamma - 1)/(\gamma - 2a - 2)$ so $\gamma > 2a + 2$.

It is underdispersed if $a < -(\mu + 1)/2$, equidispersed if $a = -(\mu + 1)/2$ or overdispersed if $a > -(\mu + 1)/2$. In particular, if $a \geq 0$ the CTP is always overdispersed.

Value

`dctp` gives the pmf, `pctp` gives the distribution function, `qctp` gives the quantile function and `rctp` generates random values.

If $a = 0$ the probability mass function, distribution function, quantile function and random generation function for the CBP distribution arise. If $b = 0$ the probability mass function, distribution function, quantile function and random generation function for the EBW distribution arise.

References

- Jose Rodriguez-Avi J, Conde-Sanchez A, Saez-Castillo AJ (2003). "A new class of discrete distributions with complex parameters." *Stat. Pap.*, **44**, 67–88. doi:10.1007/s0036200201347.
- Rodriguez-Avi J, Conde-Sanchez A, Saez-Castillo AJ, Olmo-Jimenez MJ (2004). "A triparametric discrete distribution with complex parameters." *Stat. Pap.*, **45**, 81-95. doi:10.1007/BF02778271.
- Olmo-Jimenez MJ, Rodriguez-Avi J, Cueva-Lopez V (2018). "A review of the CTP distribution: a comparison with other over- and underdispersed count data models." *Journal of Statistical Computation and Simulation*, **88**(14), 2684-2706. doi:10.1080/00949655.2018.1482897.
- Cueva-Lopez V, Olmo-Jimenez MJ, Rodriguez-Avi J (2021). "An Over and Underdispersed Biparametric Extension of the Waring Distribution." *Mathematics*, **9**(170), 1-15. doi:10.3390/math9020170.

See Also

Functions for maximum-likelihood fitting of the CTP, CBP and EBW distributions: [fitctp](#), [fitebw](#) and [fitebw](#).

Examples

```
# Examples for the function dctp
dctp(3,1,2,5)
dctp(c(3,4),1,2,5)

# Examples for the function pctp
pctp(3,1,2,3)
pctp(c(3,4),1,2,3)

# Examples for the function qctp
qctp(0.5,1,2,3)
qctp(c(.8,.9),1,2,3)

# Examples for the function rctp
rctp(10,1,1,3)
```

Description

Probability mass function, distribution function, quantile function and random generation for the Extended Biparametric Waring (EBW) distribution with parameters α and γ (or ρ).

Usage

```
debw(x, alpha, gamma, rho)
pebw(q, alpha, gamma, rho, lower.tail = TRUE)
qebw(p, alpha, gamma, rho, lower.tail = TRUE)
rebw(n, alpha, gamma, rho, lower.tail = TRUE)
```

Arguments

x	vector of (non-negative integer) quantiles.
alpha	parameter alpha (real)
gamma	parameter γ (positive)
rho	parameter rho (positive)
q	vector of quantiles.
lower.tail	if TRUE (default), probabilities are $P(X \leq x)$, otherwise, $P(X > x)$.
p	vector of probabilities.
n	number of observations. If $\text{length}(n) > 1$, the length is taken to be the number required.

Details

The EBW distribution with parameters α and γ has pmf

$$f(x|\alpha, \gamma) = C\Gamma(\alpha + x)^2 / (\Gamma(\gamma + x)x!), x = 0, 1, 2, \dots$$

where $\Gamma(\cdot)$ is the gamma function and

$$C = \Gamma(\gamma - \alpha^2) / (\Gamma(\alpha)^2\Gamma(\gamma - 2\alpha))$$

the normalizing constant.

There is an alternative parametrization in terms of α and $\rho = \gamma - 2\alpha > 0$ when $\alpha > 0$. So, introduce only α and γ or α and ρ , depending on the parametrization you wish to use.

The mean and the variance of the EBW distribution are $E(X) = \mu = \alpha^2/(\gamma - 2\alpha - 1)$ and $Var(X) = \mu(\mu + \gamma - 1)/(\gamma - 2\alpha - 2)$ so $\gamma > 2\alpha + 2$.

It is underdispersed if $\alpha < -(\mu + 1)/2$, equidispersed if $\alpha = -(\mu + 1)/2$ or overdispersed if $\alpha > -(\mu + 1)/2$. In particular, if $\alpha \geq -0.5$ the EBW is overdispersed, whereas if $\alpha < -1$ the EBW is underdispersed. In the case $-1 < \alpha \leq -0.5$, the EBW may be under-, equi- or overdispersed depending on the value of γ .

Value

debw gives the pmf, pebw gives the distribution function, qebw gives the quantile function and rebw generates random values.

If $\alpha > 0$ the probability mass function, distribution function, quantile function and random generation function for the UGW(α, α, ρ) distribution arise.

If $\alpha < 0$ the probability mass function, distribution function, quantile function and random generation function for the CTP($\alpha, 0, \gamma$) distribution arise.

References

- Jose Rodriguez-Avi J, Conde-Sanchez A, Saez-Castillo AJ (2003). "A new class of discrete distributions with complex parameters." *Stat. Pap.*, **44**, 67–88. doi:10.1007/s0036200201347.
- Rodriguez-Avi J, Conde-Sanchez A, Saez-Castillo AJ, Olmo-Jimenez MJ (2004). "A triparametric discrete distribution with complex parameters." *Stat. Pap.*, **45**, 81-95. doi:10.1007/BF02778271.
- Olmo-Jimenez MJ, Rodriguez-Avi J, Cueva-Lopez V (2018). "A review of the CTP distribution: a comparison with other over- and underdispersed count data models." *Journal of Statistical Computation and Simulation*, **88**(14), 2684-2706. doi:10.1080/00949655.2018.1482897.

See Also

Functions for maximum-likelihood fitting of the CTP and CBP distributions: [fitctp](#) and [fitcbp](#).

Examples

```
# Examples for the function dctp
debw(3,1,rho=5)
debw(c(3,4),2,rho=5)

# Examples for the function pebw
pebw(3,2,rho=5)
pebw(c(3,4),2,rho=5)

# Examples for the function qebw
qebw(0.5,-2.1,gamma=0.1)
qebw(c(.8,.9),-2.1,gamma=0.1)
qebw(0.5,2,rho=5)
qebw(c(.8,.9),2,rho=5)

# Examples for the function rebw
rebw(10,2,rho=5)
rebw(10,-2.1,gamma=5)
```


fitcbp

*Maximum-likelihood fitting of the CBP distribution***Description**

Maximum-likelihood fitting of the Complex Biparametric Pearson (CBP) distribution with parameters b and γ . Generic methods are `print`, `summary`, `coef`, `logLik`, `AIC`, `BIC` and `plot`.

Usage

```
fitcbp(x, bstart = NULL, gammastart = NULL, method = "L-BFGS-B", control = list(), ...)
```

Arguments

<code>x</code>	A numeric vector of length at least one containing only finite values.
<code>bstart</code>	A starting value for the parameter b ; by default <code>NULL</code> .
<code>gammastart</code>	A starting value for the parameter $\gamma > 0$; by default <code>NULL</code> .
<code>method</code>	The method to be used in fitting the model. See 'Details'.
<code>control</code>	A list of parameters for controlling the fitting process.
<code>...</code>	Additional parameters.

Details

If the starting values of the parameters b and γ are omitted (default option), they are computing by the method of moments (if possible; otherwise they must be entered).

The default method is "L-BFGS-B" (see details in `optim` function), but non-linear minimization (`nlm`) or those included in the `optim` function ("Nelder-Mead", "BFGS", "CG" and "SANN") may be used.

Standard error (SE) estimates for b and γ are provided by the default method; otherwise, SE for γ_0 where $\gamma = \exp(\gamma_0)$ is computed.

Value

An object of class 'fitCBP' is a list containing the following components:

- `n`, the number of observations,
- `initialValues`, a vector with the starting values used,
- `coefficients`, the parameter ML estimates of the CTP distribution,
- `se`, a vector of the standard error estimates,
- `hessian`, a symmetric matrix giving an estimate of the Hessian at the solution found in the optimization of the log-likelihood function,
- `cov`, an estimate of the covariance matrix of the model coefficients,
- `corr`, an estimate of the correlation matrix of the model estimates,

- `loglik`, the maximized log-likelihood,
- `aic`, Akaike Information Criterion, minus twice the maximized log-likelihood plus twice the number of parameters,
- `bic`, Bayesian Information Criterion, minus twice the maximized log-likelihood plus twice the number of parameters,
- `code`, a code that indicates successful convergence of the fitter function used (see `nlm` and `optim` helps),
- `converged`, logical value that indicates if the optimization algorithms successful,
- `method`, the name of the fitter function used.

Generic functions:

- `print`: The print of a 'fitCBP' object shows the ML parameter estimates and their standard errors in parenthesis.
- `summary`: The summary provides the ML parameter estimates, their standard errors and the statistic and p-value of the Wald test to check if the parameters are significant. This summary also shows the loglikelihood, AIC and BIC values, as well as the results for the chi-squared goodness-of-fit test and the Kolmogorov-Smirnov test for discrete variables. Finally, the correlation matrix between parameter estimates appears.
- `coef`: It extracts the fitted coefficients from a 'fitCBP' object.
- `logLik`: It extracts the estimated log-likelihood from a 'fitCBP' object.
- `AIC`: It extracts the value of the Akaike Information Criterion from a 'fitCBP' object.
- `BIC`: It extracts the value of the Bayesian Information Criterion from a 'fitCBP' object.
- `plot`: It shows the plot of a 'fitCBP' object. Observed and theoretical probabilities, empirical and theoretical cumulative distribution functions or empirical cumulative probabilities against theoretical cumulative probabilities are the three plot types.

References

Jose Rodriguez-Avi J, Conde-Sanchez A, Saez-Castillo AJ (2003). "A new class of discrete distributions with complex parameters." *Stat. Pap.*, **44**, 67–88. doi:10.1007/s0036200201347.

See Also

Plot of observed and theoretical frequencies for a CBP fit: [plot.fitCBP](#)

Maximum-likelihood fitting for the CTP distribution: [fitctp](#).

Maximum-likelihood fitting for the EBW distribution: [fitebw](#).

Examples

```
set.seed(123)
x <- rcbp(500, 1.75, 3.5)
fitcbp(x)
summary(fitcbp(x, bstart = 1.1, gammastart = 3))
```

fitctp	<i>Maximum-likelihood fitting of the CTP distribution</i>
--------	---

Description

Maximum-likelihood fitting of the Complex Triparametric Pearson (CTP) distribution with parameters a , b and γ . Generic methods are `print`, `summary`, `coef`, `logLik`, `AIC`, `BIC` and `plot`.

Usage

```
fitctp(x, astart = NULL, bstart = NULL, gammastart = NULL,
       method = "L-BFGS-B", control = list(), ...)
```

Arguments

<code>x</code>	A numeric vector of length at least one containing only finite values.
<code>astart</code>	A starting value for the parameter $a > 0$; by default <code>NULL</code> .
<code>bstart</code>	A starting value for the parameter b ; by default <code>NULL</code> .
<code>gammastart</code>	A starting value for the parameter $\gamma > \max(0, 2a)$; by default <code>NULL</code> .
<code>method</code>	The method to be used in fitting the model. See 'Details'.
<code>control</code>	A list of parameters for controlling the fitting process.
<code>...</code>	Additional parameters.

Details

If the starting values of the parameters a , b and γ are omitted (default option), they are computed by the method of moments (if possible; otherwise they must be entered).

The default method is "L-BFGS-B" (see details in `optim` function), but non-linear minimization (`nlm`) or those included in the `optim` function ("Nelder-Mead", "BFGS", "CG" and "SANN") may be used.

Standard error (SE) estimates for a , b and γ are provided by the default method; otherwise, SE for γ_0 where $\gamma = \exp(\gamma_0)$ is computed.

Value

An object of class 'fitCTP' is a list containing the following components:

- `n`, the number of observations,
- `initialValues`, a vector with the starting values used,
- `coefficients`, the parameter ML estimates of the CTP distribution,
- `se`, a vector of the standard error estimates,
- `hessian`, a symmetric matrix giving an estimate of the Hessian at the solution found in the optimization of the log-likelihood function,

- `cov`, an estimate of the covariance matrix of the model coefficients,
- `corr`, an estimate of the correlation matrix of the model estimates,
- `loglik`, the maximized log-likelihood,
- `aic`, Akaike Information Criterion, minus twice the maximized log-likelihood plus twice the number of parameters,
- `bic`, Bayesian Information Criterion, minus twice the maximized log-likelihood plus twice the number of parameters,
- `code`, a code that indicates successful convergence of the fitter function used (see `nlm` and `optim` helps),
- `converged`, logical value that indicates if the optimization algorithms successful,
- `method`, the name of the fitter function used.

Generic functions:

- `print`: The print of a 'fitCTP' object shows the ML parameter estimates and their standard errors in parenthesis.
- `summary`: The summary provides the ML parameter estimates, their standard errors and the statistic and p-value of the Wald test to check if the parameters are significant. This summary also shows the loglikelihood, AIC and BIC values, as well as the results for the chi-squared goodness-of-fit test and the Kolmogorov-Smirnov test for discrete variables. Finally, the correlation matrix between parameter estimates appears.
- `coef`: It extracts the fitted coefficients from a 'fitCTP' object.
- `logLik`: It extracts the estimated log-likelihood from a 'fitCTP' object.
- `AIC`: It extracts the value of the Akaike Information Criterion from a 'fitCTP' object.
- `BIC`: It extracts the value of the Bayesian Information Criterion from a 'fitCTP' object.
- `plot`: It shows the plot of a 'fitCTP' object. Observed and theoretical probabilities, empirical and theoretical cumulative distribution functions or empirical cumulative probabilities against theoretical cumulative probabilities are the three plot types.

References

- Rodriguez-Avi J, Conde-Sanchez A, Saez-Castillo AJ, Olmo-Jimenez MJ (2004). "A triparametric discrete distribution with complex parameters." *Stat. Pap.*, **45**, 81-95. doi:10.1007/BF02778271.
- Olmo-Jimenez MJ, Rodriguez-Avi J, Cueva-Lopez V (2018). "A review of the CTP distribution: a comparison with other over- and underdispersed count data models." *Journal of Statistical Computation and Simulation*, **88**(14), 2684-2706. doi:10.1080/00949655.2018.1482897.

See Also

- Plot of observed and theoretical frequencies for a CTP fit: [plot.fitCTP](#)
- Maximum-likelihood fitting for the CBP distribution: [fitcbp](#).
- Maximum-likelihood fitting for the EBW distribution: [fitebw](#).

Examples

```
set.seed(123)
x <- rctp(500, -0.5, 1, 2)
fitctp(x)
summary(fitctp(x, astart = 1, bstart = 1.1, gammastart = 3))
```

fitebw

*Maximum-likelihood fitting of the EBW distribution***Description**

Maximum-likelihood fitting of the Extended Biparametric Waring (EBW) distribution with parameters α , ρ and γ . Generic methods are `print`, `summary`, `coef`, `logLik`, `AIC`, `BIC` and `plot`. The method to be used in fitting the model is "L-BFGS-B" which allows constraints for each variable (see details in `optim` function).

Usage

```
fitebw(x, alphastart = NULL, rhostart = NULL, gammastart = NULL,
       method = "L-BFGS-B", control = list(),...)
```

Arguments

<code>x</code>	A numeric vector of length at least one containing only finite values.
<code>alphastart</code>	A starting value for the parameter α ; by default NULL.
<code>rhostart</code>	A starting value for the parameter $\rho > 0$; by default NULL.
<code>gammastart</code>	A starting value for the parameter $\gamma > \max(0, 2\alpha)$; by default NULL.
<code>method</code>	The method to be used in fitting the model. The default method is "L-BFGS-B" (<code>optim</code>).
<code>control</code>	A list of parameters for controlling the fitting process.
<code>...</code>	Additional parameters.

Details

If the starting value for α is positive, the parameterization (α, ρ) is used; otherwise, the parameterization (α, γ) is used.

If the starting values of the parameters α , γ or ρ are omitted (default option), they are computed by the method of moments (if possible; otherwise they must be entered).

The default method is "L-BFGS-B" (see details in `optim` function), but non-linear minimization (`nlm`) or those included in the `optim` function ("Nelder-Mead", "BFGS", "CG" and "SANN") may be used.

Standard error (SE) estimates for α , γ or ρ are provided by the default method; otherwise, SE for α_0 and γ_0 where $\alpha = -\exp(\alpha_0)$ and $\gamma = \exp(\gamma_0)$ (or for α_0 and ρ_0 where $\alpha = \exp(\alpha_0)$ and $\rho = \exp(\rho_0)$) are computed.

Value

An object of class 'fitEBW' is a list containing the following components:

- n, the number of observations,
- initialValues, a vector with the starting values used,
- coefficients, the parameter ML estimates of the CTP distribution,
- se, a vector of the standard error estimates,
- hessian, a symmetric matrix giving an estimate of the Hessian at the solution found in the optimization of the log-likelihood function,
- cov, an estimate of the covariance matrix of the model coefficients,
- corr, an estimate of the correlation matrix of the model estimates,
- loglik, the maximized log-likelihood,
- aic, Akaike Information Criterion, minus twice the maximized log-likelihood plus twice the number of parameters,
- bic, Bayesian Information Criterion, minus twice the maximized log-likelihood plus twice the number of parameters,
- code, a code that indicates successful convergence of the fitter function used (see nlm and optim helps),
- converged, logical value that indicates if the optimization algorithms successful.
- method, the name of the fitter function used.

Generic functions:

- print: The print of a 'fitEBW' object shows the ML parameter estimates and their standard errors in parenthesis.
- summary: The summary provides the ML parameter estimates, their standard errors and the statistic and p-value of the Wald test to check if the parameters are significant. This summary also shows the loglikelihood, AIC and BIC values, as well as the results for the chi-squared goodness-of-fit test and the Kolmogorov-Smirnov test for discrete variables. Finally, the correlation matrix between parameter estimates appears.
- coef: It extracts the fitted coefficients from a 'fitEBW' object.
- logLik: It extracts the estimated log-likelihood from a 'fitEBW' object.
- AIC: It extracts the value of the Akaike Information Criterion from a 'fitEBW' object.
- BIC: It extracts the value of the Bayesian Information Criterion from a 'fitEBW' object.
- plot: It shows the plot of a 'fitEBW' object. Observed and theoretical probabilities, empirical and theoretical cumulative distribution functions or empirical cumulative probabilities against theoretical cumulative probabilities are the three plot types.

References

Cueva-Lopez V, Olmo-Jimenez MJ, Rodriguez-Avi J (2021). "An Over and Underdispersed Biparametric Extension of the Waring Distribution." *Mathematics*, **9**(170), 1-15. doi:10.3390/math9020170.

See Also

Plot of observed and theoretical frequencies for a EBW fit: [plot.fitEBW](#)

Maximum-likelihood fitting for the CTP distribution: [fitctp](#).

Maximum-likelihood fitting for the CBP distribution: [fitcbp](#).

Examples

```
set.seed(123)
x <- rebw(500, 2, rho = 5)
fitebw(x)
summary(fitebw(x, alphastart = 1, rhostart = 5))
```

plot.fitCBP

Plot of observed and theoretical frequencies for a CBP fit

Description

Plot of observed and theoretical frequencies for a CBP fit

Usage

```
## S3 method for class 'fitCBP'
plot(x, plty = "FREQ", maxValue = NULL, ...)
```

Arguments

x	An object of class 'fitCBP'
plty	Plot type to be shown. Default is "FREQ" which shows the observed and theoretical frequencies for each value of the variable; "CDF" and "PP" are also available for plotting the empirical and theoretical cumulative distribution functions or the theoretical cumulative probabilities against the empirical cumulative probabilities, respectively.
maxValue	maxValue you want to appear in the plot
...	Additional parameters.

Examples

```
set.seed(123)
x <- rcbp(500, 1.75, 3.5)
fit <- fitcbp(x)
plot(fit)
plot(fit, plty = "CDF")
plot(fit, plty = "PP")
```

plot.fitCTP *Plot of observed and theoretical frequencies for a CTP fit*

Description

Plot of observed and theoretical frequencies for a CTP fit

Usage

```
## S3 method for class 'fitCTP'
plot(x, plty = "FREQ", maxValue = NULL, ...)
```

Arguments

x	An object of class 'fitCTP'
plty	Plot type to be shown. Default is "FREQ" which shows the observed and theoretical frequencies for each value of the variable; "CDF" and "PP" are also available for plotting the empirical and theoretical cumulative distribution functions or the theoretical cumulative probabilities against the empirical cumulative probabilities, respectively.
maxValue	maxValue you want to appear in the plot
...	Additional parameters.

Examples

```
set.seed(123)
x <- rctp(500, -0.5, 1, 2)
fit <- fitctp(x)
plot(fit)
plot(fit, plty = "CDF")
plot(fit, plty = "PP")
```

plot.fitEBW *Plot of observed and theoretical frequencies for a EBW fit*

Description

Plot of observed and theoretical frequencies for a EBW fit

Usage

```
## S3 method for class 'fitEBW'
plot(x, plty = "FREQ", maxValue = NULL, ...)
```


Arguments

<code>x</code>	An object of class 'fitEBW'
<code>plty</code>	Plot type to be shown. Default is "FREQ" which shows the observed and theoretical frequencies for each value of the variable; "CDF" and "PP" are also available for plotting the empirical and theoretical cumulative distribution functions or the theoretical cumulative probabilities against the empirical cumulative probabilities, respectively.
<code>maxValue</code>	maxValue you want to appear in the plot
<code>...</code>	Additional parameters.

Examples

```
set.seed(123)
x <- rebw(500, -0.25, 1)
fit <- fitebw(x)
plot(fit)
plot(fit, plty = "CDF")
plot(fit, plty = "PP")
```

varcomp

Variance decomposition for a EBW fit

Description

One of the main drawbacks of the Univariate Generalized Waring (UGW) distribution with parameters a , k and ρ is that the first two parameters are interchangeable, so it is not possible to distinguish the variance components 'liability' and 'proneness' without additional information. To solve this problem, an EBW distribution (where these components are uniquely identifiable) can be used since, given a UGW distribution, there always exists an EBW close to it.

Usage

```
varcomp(object, ...)
```

Arguments

<code>object</code>	An object of class 'fitEBW'
<code>...</code>	Additional parameters.

Value

A data frame with the variance components (randomness, liability and proneness) in absolute and relative terms.

Examples

```
set.seed(123)
x <- rebw(500, 2, rho = 5)
fit <- fitebw(x, alphastart = 1, rhostart = 5)
varcomp(fit)
```

Index

cbp, 2
chisq.test2, 4
ctp, 5

dcbp (cbp), 2
dctp, 3
dctp (ctp), 5
debw (ebw), 7

ebw, 7

fitcbp, 3, 6, 8, 9, 12, 15
fitctp, 6, 8, 10, 11, 15
fitebw, 6, 10, 12, 13

nlm, 9, 11, 13

optim, 9, 11, 13

pcbp (cbp), 2
pctp, 3
pctp (ctp), 5
pebw (ebw), 7
plot.fitCBP, 10, 15
plot.fitCTP, 12, 16
plot.fitEBW, 15, 16

qcbp (cbp), 2
qctp, 3
qctp (ctp), 5
qebw (ebw), 7

rcbp (cbp), 2
rctp, 3
rctp (ctp), 5
rebw (ebw), 7

varcomp, 17